arXiv:1701.07800v1 [math.CA] 26 Jan 2017

A MULTILINEAR REVERSE HÖLDER INEQUALITY WITH APPLICATIONS TO MULTILINEAR WEIGHTED NORM INEQUALITIES

DAVID CRUZ-URIBE, OFS AND KABE MOEN

Abstract. In this note we prove a multilinear version of the reverse Hölder inequality in the theory of Muckenhoupt A_p weights. We give two applications of this inequality to the study of multilinear weighted norm inequalities. First, we prove a structure theorem for multilinear $A_{\vec{P}}$ weights; second, we give a new sufficient condition for multilinear, two-weight norm inequalities for the maximal operator.

1. Introduction

The purpose of this note is to prove a multilinear version of the reverse Hölder inequality in the theory of the Muckenhoupt A_p classes. We briefly recall the definitions; for complete details, see [5]. For $1 < p < \infty$, a weight $w \in A_p$ if

$$[w]_{A_p} = \sup_Q \left(\fint_Q w\,dx\right)\left(\fint_Q w^{1-p'}\,dx\right)^{p-1} < \infty;$$

here and below, the supremum is taken over all cubes in $\mathbb{R}^n$ with sides parallel to the coordinate axes. We denote the union of all the A_p classes by A_∞. If $w \in A_\infty$, then it satisfies the reverse Hölder inequality: there exists $s > 1$ such that

$$[w]_{RH_s} = \sup_Q \left(\fint_Q w^s\,dx\right)^{\frac{1}{s}}\left(\fint_Q w\,dx\right)^{-1} < \infty.$$

Our main result extends this condition to m-tuples of A_∞ weights.

Theorem 1.1. *Given $1 < s_1, \dots, s_m < \infty$ such that $\sum \frac{1}{s_i} = 1$, suppose $w_i \in RH_{s_i}$. Then for all cubes Q,*

$$\prod_{i=1}^m \left(\fint_Q w_i^{s_i}\,dx\right)^{\frac{1}{s_i}} \le C \fint_Q \prod_{i=1}^m w_i\,dx. \tag{1.1}$$

Date: 1/26/2017.

2000 *Mathematics Subject Classification.* 42B20, 42B25.

The first author is supported by NSF Grant DMS-1362425 and research funds from the Dean of the College of Arts & Sciences, the University of Alabama. The second author is supported by the Simons Foundation.

The proof requires the rescaling properties of Muckenhoupt weights from [4, Theorems 2.1, 2.2] which we state here as a lemma. (Also see [9].)

Lemma 1.2. *Given a weight w:*

(a) *$w \in RH_s$, $1 < s < \infty$, if and only if $w^s \in A_\infty$;*
(b) *$w \in A_p \cap RH_s$, $1 < p$, $s < \infty$ if and only if $w^s \in A_q$, $q = s(p-1)+1$.*

Proof. By Lemma 1.2, for each i, since $w_i \in RH_{s_i}$, $w_i^{s_i} \in A_\infty$. Moreover, there exists $0 < r < 1$ such that $w_i^{rs_i} \in A_2 \cap RH_{\frac{1}{r}}$; since the A_p classes are nested ($A_p \subset A_q$ if $p < q$) we can find a single value of r that works for all i. Therefore, if we apply the reverse Hölder inequality, the A_2 condition, multilinear Hölder's inequality with exponents s_i, and then Hölder's inequality twice more, we get that for every cube Q,

$$\begin{aligned}\prod_{i=1}^m \left(\fint_Q w_i^{s_i}\,dx\right)^{\frac{1}{s_i}} &\le C\prod_{i=1}^m \left(\fint_Q w_i^{rs_i}\,dx\right)^{\frac{1}{rs_i}} \le C\prod_{i=1}^m \left(\fint_Q w_i^{-rs_i}\,dx\right)^{-\frac{1}{rs_i}}\\ &\le C\left(\fint_Q \prod_{i=1}^m w_i^{-r}\,dx\right)^{-\frac{1}{r}} \le C\left(\fint_Q \prod_{i=1}^m w_i^{r}\,dx\right)^{\frac{1}{r}} \le C\fint_Q \prod_{i=1}^m w_i\,dx.\end{aligned}$$

□

Remark 1.3. In the bilinear case, a version of this result was proved in [4, Theorem 2.6]. However, our proof here is considerably simpler.

Theorem 1.6 has two corollaries. The first when $p = 2$ is part of the folklore of harmonic analysis, but we have not been able to find a proof in the literature. We include this result since it bears a resemblance to Theorem 1.6 below.

Corollary 1.4. *For $1 < p < \infty$, if a weight w is such that w, $w^{1-p'} \in A_\infty$, then $w \in A_p$.*

Proof. Let $w_1 = w^{\frac{1}{p}}$, $w_2 = w^{-\frac{1}{p}}$, and $s_1 = p$, $s_2 = p'$. Then by Lemma 1.2, $w_i \in RH_{s_i}$, and inequality (1.1) becomes the A_p condition. □

Our second corollary is a version of Theorem 1.1 that is particularly useful for applications to multilinear weights.

Corollary 1.5. *Let $1 < p_1, \dots, p_m < \infty$, $0 < p < \infty$ be such that $\sum_{i=1}^m \frac{1}{p_i} = \frac{1}{p}$, and let $\vec{w} = (w_1, \dots, w_m)$ be a vector of weights. If $w_i \in A_\infty$, $1 \le i \le m$, then for all cubes Q,*

$$\prod_{i=1}^m \left(\fint_Q w_i\,dx\right)^{\frac{p}{p_i}} \le C\fint_Q \prod_{i=1}^m w_i^{\frac{p}{p_i}}\,dx.$$

Proof. By Lemma 1.2, for each i, since $w_i \in A_\infty$ we have $w_i^{\frac{p}{p_i}} \in RH_{\frac{p_i}{p}}$. The desired inequality now follows from Theorem 1.1 with $s_i = \frac{p_i}{p}$. □

We now give two applications of Theorem 1.1 (or more properly, Corollary 1.5) to the theory of multilinear weights. We first consider the structure of multilinear $A_{\vec{p}}$ weights. To put our result into context, we briefly sketch the history of multilinear weighted norm inequalities. A multilinear Calderón-Zygmund singular integral operator T is an m-linear operator with kernel K such that

$$T(\vec{f})(x) = \int_{\mathbb{R}^{mn}} K(x, y_1, \dots, y_m) \prod_{i=1}^{m} f_i(y_i)\, dy_1 \dots dy_m,$$

where $\vec{f} = (f_i, \dots, f_m)$, $f_i \in C_c^\infty(\mathbb{R}^n)$ and $x \notin \bigcap_{i=1}^m \operatorname{supp}(f_i)$, and the kernel satisfies certain regularity estimates. (Precise definitions are not necessary for our purposes; see Grafakos and Torres [8] for details and further references.) These operators satisfy the following L^p space estimates: if $1 < p_1, \dots, p_m < \infty$ and $\frac{1}{p} = \sum \frac{1}{p_i}$, then

$$\|T(\vec{f})\|_{L^p} \le C \prod_{i=1}^{m} \|f_i\|_{L^{p_i}}.$$

Weighted norm inequalities for multilinear singular integral operators were first considered by Grafakos and Torres [7]. They showed that if $p_0 = \min(p_1, \dots, p_m)$ and $w \in A_{p_0}$, then

$$\|T(\vec{f})\|_{L^p(w)} \le C \prod_{i=1}^{m} \|f_i\|_{L^{p_i}(w)}.$$

This result was generalized by Grafakos and Martell [6], who showed that if $w_i \in A_{p_i}$ and $w = \prod w_i^{\frac{p}{p_i}}$, then

$$\|T(\vec{f})\|_{L^p(w)} \le C \prod_{i=1}^{m} \|f_i\|_{L^{p_i}(w_i)}. \tag{1.2}$$

The optimal class of weights for which (1.2) holds was found by Lerner, *et al.* [10]. Given $\vec{p} = (p_1, \dots, p_m)$, $\vec{w} = (w_1, \dots, w_m)$ and $w = \prod w_i^{\frac{p}{p_i}}$, $\vec{w} \in A_{\vec{p}}$ if

$$[\vec{w}]_{A_{\vec{p}}} = \sup_Q \left(\fint_Q w\, dx \right)^{\frac{1}{p}} \prod_{i=1}^{m} \left(\fint_Q w_i^{1-p_i'}\, dx \right)^{\frac{1}{p_i'}} < \infty.$$

It follows at once by Hölder's inequality that if $w_i \in A_{p_i}$, then $\vec{w} \in A_{\vec{p}}$. Such weights are referred to as product $A_{\vec{p}}$ weights.[1] However, in [10] they constructed examples of weights $\vec{w} \in A_{\vec{p}}$ such that the weights $w_i \notin L^1_{\mathrm{Loc}}$ and so not in A_{p_i}.

Here we prove that if we impose some additional regularity on the weights w_i, then the weight $\vec{w}$ is a product weight.

Theorem 1.6. *Given $1 < p_1, \dots, p_m < \infty$ and $\frac{1}{p} = \sum \frac{1}{p_i}$, suppose $\vec{w} = (w_1, \dots, w_m) \in A_{\vec{p}}$. If $w_i \in A_\infty$, $1 \le i \le m$, then $w_i \in A_{p_i}$.*

Proof of Theorem 1.6. By Hölder's inequality, for every i and every cube Q,

$$1 \le \left(\fint_Q w_i\,dx\right)^{\frac{p}{p_i}} \left(\fint_Q w_i^{1-p_i'}\,dx\right)^{\frac{p}{p_i'}}.$$

Thus, if we fix a value of i,

$$\begin{aligned}\left[\fint_Q w_i\,dx \left(\fint_Q w_i^{1-p_i'}\,dx\right)^{p_i-1}\right]^{\frac{p}{p_i'}} &= \left(\fint_Q w_i\,dx\right)^{\frac{p}{p_i}} \left(\fint_Q w_i^{1-p_i'}\,dx\right)^{\frac{p}{p_i'}} \\ &\le \prod_{i=1}^m \left(\fint_Q w_i\,dx\right)^{\frac{p}{p_i}} \left(\fint_Q w_i^{1-p_i'}\,dx\right)^{\frac{p}{p_i'}}.\end{aligned}$$

By Corollary 1.5,

$$\begin{aligned}&\prod_{i=1}^m \left(\fint_Q w_i\,dx\right)^{\frac{p}{p_i}} \left(\fint_Q w_i^{1-p_i'}\,dx\right)^{\frac{p}{p_i'}} \\ &\qquad\le C\left(\fint_Q \prod_{i=1}^m w_i^{\frac{p}{p_i}}\,dx\right) \prod_{i=1}^m \left(\fint_Q w_i^{1-p_i'}\,dx\right)^{\frac{p}{p_i'}} \le C[\vec{w}]_{A_{\vec{p}}}.\end{aligned}$$

If we combine these two inequalities, we see that $w_i \in A_{p_i}$. □

It is tempting to speculate that the condition $w_i \in A_\infty$ in Theorem 1.6 can be weakened to $w_i \in L^1_{\mathrm{Loc}}$. This, however, is false, and we construct a counter-example. We note in passing that our example also gives an example of a weight $\vec{w} \in A_{\vec{p}}$ that is not a product weight; our construction is somewhat simpler than that given in [10, Remark 7.2].

For simplicity we consider the bilinear case $m = 2$ on the real line and we will use the characterization $\vec{w} \in A_{\vec{p}}$ if and only if $w \in A_{2p}$ and $\sigma_i = w^{1-p_i'} \in A_{2p_i'}$ (see [10,

[1] If the weights w_i are taken as prior, then this name makes sense. If the weight w is taken as prior, then it makes more sense to refer to these weights as "factored" $A_{\vec{p}}$ weights. However, "product weights" has become the standard terminology.

Theorem 3.6]). Let

$$w_1(x) = \frac{\chi_{[-1,1]}(x)}{|x|\log(e/|x|)^2} + \chi_{[-1,1]^c}(x) \quad \text{and} \quad w_2(x) = 1.$$

First, we have that $w = w_1^{\frac{p}{p_1}}$ belongs to $A_1 \subset A_{2p}$. To see this, note that for $0 < a < 1$ and $b > 0$, if we define

$$v(x) = \frac{\chi_{[-1,1]}(x)}{|x|^a \log(e/|x|)^b} + \chi_{[-1,1]^c}(x),$$

then $v \in A_1$. To show this, it suffices to check the A_1 condition on intervals $[0,t]$, $0 < t < 1$ (cf. [3]), but in this case it is straightforward to show that

$$\int_0^t \frac{ds}{s^a \log(e/s)^b} \approx \frac{1}{t^{a-1}\log(e/t)^b}. \tag{1.3}$$

(Just compare the integrand to the derivative of the righthand side.) Since $p/p_1 < 1$ we have that $w \in A_1$. Next we will show that the weight $\sigma_1 = w_1^{1-p_1'}$ belongs to $A_{2p_1'}$. To see this, consider the dual weight (dual in the class $A_{(2p_1)'}$) $\sigma_1^{1-(2p_1')'}$ which is given by

$$\sigma_1(x)^{1-(2p_1')'} = w_1(x)^{(p_1'-1)((2p_1')'-1)} = w_1(x)^{\frac{p_1'-1}{2p_1'-1}}.$$

Again, since $\frac{p_1'-1}{2p_1'-1} < 1$ the calculation (1.3) shows that $\sigma_1^{1-(2p_1')'} \in A_1 \subset A_{(2p_1')'}$ which implies $\sigma_1 \in A_{2p_1'}$. This shows $\vec{w} \in A_{\vec{p}}$. However, $w_i \notin A_{p_i}$: since $w_1 \in L^1_{\mathrm{Loc}} \backslash \bigcup_{p>1} L^p_{\mathrm{Loc}}$ it cannot satisfy a reverse Hölder inequality and so is not in A_∞.

Remark 1.7. It would be interesting to have an intrinsic characterization of those $A_{\vec{p}}$ weights that can be factored as product weights.

Our second application of Corollary 1.5 is to two-weight multilinear norm inequalities. In [10], the authors introduced a new multilinear maximal function as a tool to prove the estimates (1.2):

$$\mathcal{M}(\vec{f})(x) = \sup_{Q \ni x} \prod_{i=1}^m \left(\fint_Q |f(y_i)|\, dy_i \right).$$

They proved that

$$\|\mathcal{M}(\vec{f})\|_{L^p(w)} \leq C \prod_{i=1}^m \|f_i\|_{L^{p_i}(w_i)}$$

if and only if $\vec{w} \in A_{\vec{p}}$. This result led naturally to the question of proving two-weight inequalities for $\mathcal{M}$: more precisely, inequalities of the form

$$\|\mathcal{M}(\vec{f\sigma})\|_{L^p(u)} \leq C \prod_{i=1}^{m} \|f_i\|_{L^{p_i}(\sigma_i)}, \tag{1.4}$$

where $\vec{f\sigma} = (f_1\sigma_1, \dots, f_m\sigma_m)$. Chen and Damián [2] defined a multilinear analog of the Sawyer testing condition for the Hardy-Littlewood maximal operator: $(u, \vec{\sigma}) \in S_{\vec{p}}$ if there exists a constant C such that for all cubes Q,

$$\left(\int_Q \mathcal{M}(\vec{\sigma}\chi_Q)^p u\,dx\right)^{\frac{1}{p}} \leq C \prod_{i=1}^{m} \sigma_i(Q)^{\frac{1}{p_i}}.$$

The $S_{\vec{p}}$ condition is necessary for the two weight norm inequality (1.4), but it is not known if it is also sufficient. Chen and Damián [2, Theorem 3.2] proved that it is sufficient if they also assume that the weights satisfy a multilinear reverse Hölder condition: for all cubes Q,

$$\prod_{i=1}^{m} \left(\fint_Q \sigma_i\,dx\right)^{\frac{p}{p_i}} \leq C \fint_Q \prod_{i=1}^{m} \sigma_i^{\frac{p}{p_i}}\,dx. \tag{1.5}$$

Therefore, by Corollary 1.5, if we assume the weights σ_i are in A_∞, then we get that the condition $S_{\vec{p}}$ is sufficient the two weight norm inequality (1.4).

Corollary 1.8. *Suppose $u, \sigma_1, \dots, \sigma_m$ are weights, $1 < p_1, \dots, p_m < \infty$, and p is defined by $\frac{1}{p} = \sum_{i=1}^{m} \frac{1}{p_i}$. If $\sigma_i \in A_\infty$ and $(u, \vec{\sigma}) \in S_{\vec{p}}$, then the two-weight norm inequality (1.4) holds.*

Remark 1.9. It is an open question to characterize the m-tuples of weights for which (1.5) holds. A version of this question was posed in [4, Theorem 2.7]; they showed that when $m = 2$, if (1.1) holds and $w_1 \in RH_{s_1}$, then $w_2 \in RH_{s_2}$.

Remark 1.10. For an additional application of Corollary 1.5, see [1], where it is used to prove necessary conditions on b for the bilinear commutators $[b, T]_i$ to satisfy weighted norm inequalities, where T is a bilinear Calderón-Zygmund singular integral or a bilinear fractional integral operator.

References

[1] L. Chaffee and D. Cruz-Uribe. Necessary conditions for the boundedness of linear and bilinear commutators on banach function spaces. *preprint*, 2017.

[2] W. Chen and W. Damián. Weighted estimates for the multisublinear maximal function. *Rend. Circ. Mat. Palermo (2)*, 62(3):379–391, 2013.

[3] D Cruz-Uribe. Piecewise monotonic doubling measures. *The Rocky Mountain Journal of Mathematics*, 26(2):545–583, 1996.

[4] D. Cruz-Uribe and C. J. Neugebauer. The structure of the reverse Hölder classes. *Trans. Amer. Math. Soc.*, 347(8):2941–2960, 1995.
[5] J. Duoandikoetxea. *Fourier analysis*, volume 29 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001.
[6] L. Grafakos and J. M. Martell. Extrapolation of weighted norm inequalities for multivariable operators and applications. *J. Geom. Anal.*, 14(1):19–46, 2004.
[7] L. Grafakos and R. H. Torres. Maximal operator and weighted norm inequalities for multilinear singular integrals. *Indiana Univ. Math. J.*, 51(5):1261–1276, 2002.
[8] L. Grafakos and R. H. Torres. Multilinear Calderón-Zygmund theory. *Adv. Math.*, 165(1):124–164, 2002.
[9] R. Johnson and C. J. Neugebauer. Homeomorphisms preserving A_p. *Rev. Mat. Iberoamericana*, 3(2):249–273, 1987.
[10] A. Lerner, S. Ombrosi, C. Pérez, R. H. Torres, and R. Trujillo-González. New maximal functions and multiple weights for the multilinear Calderón-Zygmund theory. *Adv. Math.*, 220(4):1222–1264, 2009.

David Cruz-Uribe, OFS, Department of Mathematics, University of Alabama, Tuscaloosa, AL 35487, USA
E-mail address: dcruzuribe@ua.edu

Kabe Moen, Department of Mathematics, University of Alabama, Tuscaloosa, AL 35487, USA
E-mail address: kabe.moen@ua.edu